

Dark Energy Problem in a Four-Fermion Interaction Model

Tomohiro Inagaki*

Information Media Center, Hiroshima University, Hiroshima 739-8521, Japan

Xinhe Meng†

*Institute of Physics, Nankai University, Tianjin, 300071, P.R.China(Post address),
Institute of Theoretical Physics, Academic Sinica, Beijing 100080, P.R.China*

Tsukasa Murata‡

Department of Physics, Hiroshima University, Hiroshima 739-8521, Japan

(Dated: February 1, 2008)

Cosmology enters precision testing period with the observational experiments advancement. We have not arrived to decipher the origin of the dominant dark energy component of our universe, although we have located some of its characters. In such a situation the possibilities of a simple four-fermion interaction model are investigated to describe dark energy characters. We explicitly calculate the vacuum expectation value of the energy-momentum tensor and discuss the results with comparison to the equation of state for dark energy requirement. It is found that the negative pressure is realized at low temperature. It is an appropriate property as a dark energy candidate. We also investigate some cosmological tests for the dark energy via its equation of state.

PACS numbers: 11.30.Qc, 98.80.Cq

I. INTRODUCTION

Cosmological problems are as old as human history, but cosmology is a relatively new science. In recent years two main observational results have revealed a profound implication in our current knowledge of the universe. These are the confirmation of the flat geometry and its accelerating expansion of our universe in the relatively recent epoch, i.e., with redshift of roughly 0.5 or less, driving by an unidentified component as coined Dark Energy by M. Turner [1].

The recent observations have dramatically changed our previous intuition. The evolution of the universe was slowing down due to gravitational attractive force and also alleviated the universe age crisis problem with respect to the oldest stars and globular clusters in our Milk Way Galaxy. Respectively, the first acoustic peak of the temperature fluctuations on the cosmic microwave background radiation (CMBR) is observed by the experiments of BOOMERANG, MAXIMA and WMAP [2]. It favors a spatially flat universe and indicates our universe dominated by about 2/3 dark energy and 1/3 matter (with mostly the named non-relativistic cold dark matter) components. The direct evidences from measurements of luminosity-redshift relations of SN Ia [3] suggest that there exists a mysterious component so called dark energy with a negative pressure. It is as large as an anti-gravity contribution to accelerate the universe expansion. A simple candidate which shows the proper-

ties of the dark energy is found in the vacuum energy or cosmological constant. However, the energy scale to the cosmological constant is about 120 orders of magnitude smaller than that of the vacuum energy at the Planck time. Some fine tuning of parameters is necessary in almost all the models of particle physics. Thus a naturalness problem exists in some simple candidates. Furthermore, recent analysis of the observational data from WMAP combined with 2dF, SDSS, DASI and the previous CMB observations[4] disfavors the simplest cosmological constant explanation.

When describing the evolution of the universe, normally we start with solving the Einstein equation. Since we know that recent observations favors the flat, homogeneous and isotropic universe in Large Scale Survey, it is a good approximation to use the energy-momentum tensor for the dominant matter or radiation mainly depends on time or temperature when we consider the global evolution of the universe. For the perfect fluid model, the energy-momentum tensor consists of the total energy density ρ and smoothly distributed pressure p by $T_{\alpha\beta} = \text{diag}(\rho, p, p, p)$.

The isotropic and homogeneous universe can be explicated by Friedmann-Robertson-Walker (FRW) metrics. In this case Einstein equation reduces to only two relations. One is Friedmann equation :

$$\frac{\dot{a}^2}{a^2} = \frac{8\pi G}{3}\rho - \frac{k}{a^2} + \frac{\Lambda}{3}, \quad (1)$$

where $a(t)$ is the scale factor in the FRW metrics. The other is the acceleration equation

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p) + \frac{\Lambda}{3}. \quad (2)$$

These equations are very successful to describe the evolution history of the universe from the "First Three Min-

*inagaki@hiroshima-u.ac.jp

†mengxh@public.tpt.tj.cn

‡murata@hiroshima-u.ac.jp

utes" so far as we have observed. Furthermore, these can accommodate the relatively recent accelerating expansion of our universe if we accept the existence of a kind of vacuum energy or the cosmological constant-like term. On the other hand we have another solution to introduce an entity endowed with the sort of dark energy [5]. Its energy density can govern the cosmological evolution in Friedmann equations of standard cosmology [1]. With the present observational data we can not finally distinguish yet whether the accelerating expansion comes from the cosmological constant or not.

The cosmological constant can be absorbed into the energy density ρ and pressure p by a re-definition in the above equations. To reconcile with the accelerating expansion of the universe we must have

$$-1 \leq \frac{p}{\rho} < -\frac{1}{3}. \quad (3)$$

The lower bound, $p/\rho = -1$, corresponds to the cosmological constant. When we consider the dark energy model that causes the accelerating expansion, the dominant substance has to satisfy this relation by its own ρ and p .

Lots of previous works are reported in concentrating on a dynamical scalar field such as quintessence, spin-essence and so on. In such investigations a slowly varying energy density has been employed to mimic an effective cosmological constant [6]. As far as we know a very little number of works have been done in the study of the fermion field as the possible dark energy candidate.

One of the interesting candidate of the dark energy is postulated in the model with a fermion field. In some strong coupling fermion models dynamical symmetry breaking takes place by the fermion and the anti-fermion condensation below a critical temperature. As is well known, the chiral symmetry for quarks is broken down dynamically in QCD. The composite operator of the fermion and the anti-fermion develops the non-vanishing vacuum expectation value. It can change behaviors of the energy density and pressure of the universe at low temperature as the relevant energy scale in the model can be adjusted accordingly. Therefore we launched our plan to make a systematic study of the strong coupling fermion models at finite temperature for a dark energy candidate.

Here we take a four-fermion interaction model as one of such prototype models and discuss the possibilities to describe the dark energy characters by using the thermal quantum field theory technique [7]. The spacetime curvature also has an important effect on the phase structure of dynamical symmetry breaking in the early stage of the universe [8, 9, 10]. However, we consider the era when the main contribution to the evaluation of the energy-momentum tensor comes from the thermal effect. We assume that the spacetime curvature and the expansion of the universe do not have a large influence to the evaluation of the energy-momentum tensor. As is known, the four-fermion interaction is non-renormalizable. We re-

gard the model is an effective theory which is stemming from a more fundamental theory at a high energy scale.

In the present paper we study a four-fermion interaction model as a dark energy candidate. By using the imaginary time formalism we calculate the energy-momentum tensor in our model at finite temperature. The EOS is obtained from the energy-momentum tensor. Behaviors of the deceleration parameter are discussed to see the dark energy property of the four-fermion interaction model. Throughout this paper we use natural units with $c = k_B = \hbar = 1$ and assume the universe is geometric flat as astrophysical observation favors.

II. ENERGY-MOMENTUM TENSOR IN A FOUR-FERMION INTERACTION MODEL

First we briefly review the definition of the energy-momentum tensor in the quantum field theory of fermion. The four-fermion interaction model is one of the simplest model where dynamical symmetry breaking takes place [11, 12]. In D -dimensional spacetime it is described by the action :

$$S = \int d^D x \det(V) \times \left[\frac{i}{2} \sum_{k=1}^N \{ \bar{\psi}_k \gamma^a V_a^\nu \nabla_\nu \psi_k - V_a^\nu (\nabla_\nu \bar{\psi}_k) \gamma^a \psi_k \} + \frac{\lambda}{2N} \left(\sum_{k=1}^N \bar{\psi}_k \psi_k \right)^2 \right], \quad (4)$$

where the λ is a coupling constant, N is the number of fermion flavors and V_a^μ is a vierbein that relates the metric $g_{\mu\nu}$ and the local Lorentz metric η_{ab} by $g_{\mu\nu} = V_\mu^a V_\nu^b \eta_{ab}$. Below we omit the flavor index k of fermion field. The action (4) has the global chiral symmetry under the transformation $\psi \rightarrow e^{i\gamma_5 \theta} \psi$. This symmetry prevent the fermion from having mass term. Applying the auxiliary field method we rewrite the action as

$$S = \int d^D x \det(V) \times \left[\frac{i}{2} \{ \bar{\psi} \gamma^a V_a^\nu \nabla_\nu \psi - V_a^\nu (\nabla_\nu \bar{\psi}) \gamma^a \psi \} - \frac{N}{2\lambda} \sigma^2 - \bar{\psi} \sigma \psi \right]. \quad (5)$$

The non-vanishing vacuum expectation value of σ denotes the existence of the fermion and anti-fermion condensation, $\langle \sigma \rangle \sim -\frac{\lambda}{N} \langle \bar{\psi} \psi \rangle \neq 0$. It is the signal for chiral symmetry breaking. If the chiral symmetry broken down dynamically, the fermion acquire a mass term proportional to $\langle \sigma \rangle$.

The classical definition of the energy-momentum ten-

sor is [13]

$$T_{\mu\nu}(x) = \frac{2}{[-g(x)]^{1/2}} \frac{\delta S}{\delta g^{\mu\nu}(x)} = \frac{V_{a(\mu}(x)}{\det[V(x)]} \frac{\delta S}{\delta V_a^{\nu)}(x)}. \quad (6)$$

Note that the labels a, b refer to the local Lorentz frame associated with the normal coordinates, while μ, ν denote the general coordinate system x^μ . In the quantum field theory the expectation value of $T_{\mu\nu}$ is defined in the sense of the path integral formalism as

$$\begin{aligned} \langle T_{\mu\nu} \rangle &= \frac{1}{Z} \int \mathcal{D}([-g]^{1/4}\Psi) \mathcal{D}([-g]^{1/4}\bar{\Psi}) \\ &\times \mathcal{D}([-g]^{1/4}\sigma) \frac{V_{a(\mu}(x)}{\det[V(x)]} \frac{\delta S}{\delta V_a^{\nu)}(x)} e^{iS_y}, \end{aligned} \quad (7)$$

This equation is rewritten as

$$\langle T_{\mu\nu} \rangle = \frac{V_{a(\mu}(x)}{\det[V(x)]} \frac{\delta}{\delta V_a^{\nu)}(x)} (-i \ln Z). \quad (8)$$

where Z is the generating functional which is defined by

$$Z = \int \mathcal{D}([-g]^{1/4}\Psi) \mathcal{D}([-g]^{1/4}\bar{\Psi}) \mathcal{D}([-g]^{1/4}\sigma) e^{iS_y}. \quad (9)$$

We would like to discuss the evolution of the energy-momentum tensor to study a fermionic dark energy model. As a first step we confine ourselves to the Minkowski spacetime, here. The Friedmann-Robertson-Walker metric for the spatial flatness is conformally related to a Minkowski line element. For the Minkowski metric the generating functional is give by

$$-i \ln Z = N \left\{ \Gamma[\sigma] + O(1/N) \right\}, \quad (10)$$

where $\Gamma[\sigma]$ is the effective action :

$$\Gamma[\sigma] = \int d^D x \left\{ \frac{-\sigma^2}{2\lambda_0} - i \text{Tr} \ln(i\gamma^\mu \partial_\mu - \sigma) \right\}. \quad (11)$$

The energy-momentum tensor (8) contains a zero-point vacuum energy and is divergent. We use the point splitting prescription to regularize the expectation value of it [13]. Substituting the Eqs.(10) and (11) into Eq.(8) we obtain

$$\langle T_{\mu\nu}(x) \rangle = \eta_{\mu\nu} \frac{N}{2\lambda_0} \sigma^2 - iN \lim_{x' \rightarrow x} \text{Tr} \gamma_{(\mu} \partial_{\nu)}^x S_F(x, x'; \sigma), \quad (12)$$

in the leading order of $1/N$ -expansion where ∂^x denote the differentiation with respect to x and $S_F(x, x'; \sigma)$ is the Feynman propagator for the free fermion with mass σ which satisfies

$$(i\gamma^\alpha \partial_\alpha - \sigma) S_F(x, x'; \sigma) = \delta^D(x, x'). \quad (13)$$

The second term in the right hand side of Eq.(12) corresponds to the expectation value of the energy-momentum

tensor of the free fermion with mass σ . Therefore the energy-momentum tensor of the four-fermion interaction model is given by the sum of the quadratic potential term of σ , i.e., the first term in the right hand side of Eq.(12), and the energy-momentum tensor of the massive free fermion in the leading order of $1/N$ -expansion. It should be noted that the fermion mass σ is a dynamical variable. We can determine the dynamically generated fermion mass by evaluating the effective potential.

III. ENERGY-MOMENTUM TENSOR AT FINITE TEMPERATURE

Next we study the thermal effect to the energy-momentum tensor. As is shown in Eq.(12), the vacuum expectation value for the energy-momentum tensor is described by the free fermion propagator. Following the imaginary time formalism, the fermion propagator at finite temperature is obtained from the one at $T = 0$ by the Wick rotation and the replacements [7, 14]

$$\int dk^0 \rightarrow \frac{1}{\beta} \sum_n, \quad (14)$$

$$k^0 \rightarrow i\omega_n, \quad (15)$$

$$\gamma^0 \rightarrow -i\gamma^4, \quad (16)$$

where $\beta = 1/T$. The discrete variable ω_n is given by

$$\omega_n \equiv i \frac{2n+1}{\beta} \pi, \quad (17)$$

according to the anti-periodic boundary condition of the fermion field. After these replacement and some calculations the energy-momentum tensor (12) at finite temperature in the leading order of the $1/N$ expansion reads

$$\begin{aligned} \langle T^{00} \rangle &= N\sigma^2/2\lambda \\ &- \frac{2N \text{tr} 1}{(4\pi)^{(D-1)/2} \Gamma(\frac{D-1}{2})} \int_0^\infty dK K^{(D-2)} \\ &\times \left[\frac{\sqrt{K^2 + \sigma^2}}{2} \tanh\left(\frac{\beta}{2} \sqrt{K^2 + \sigma^2}\right) - \sum \frac{1}{\beta} \right] \\ &- \langle 0|T^{00}|0 \rangle, \end{aligned} \quad (18)$$

and

$$\begin{aligned} \langle T^{ii} \rangle &= -N\sigma^2/2\lambda \\ &- \frac{2N \text{tr} 1}{(D-1)(4\pi)^{(D-1)/2} \Gamma(\frac{D-1}{2})} \int_0^\infty dK K^{(D-2)} \\ &\times \left[\frac{K^2}{2\sqrt{K^2 + \sigma^2}} \tanh\left(\frac{\beta}{2} \sqrt{K^2 + \sigma^2}\right) \right] \\ &- \langle 0|T^{ii}|0 \rangle, \end{aligned} \quad (19)$$

where $\text{tr} 1$ is the trace with respect to the fermion legs in an arbitrary dimensions D , $\text{tr} 1 = 2^{D/2}$. Since the zero-point of energy-momentum tensor is unknown parameter




FIG. 1: Behavior of the expectation value of σ for $\lambda = -20\sigma_0^{2-D}$ as a function of T . σ_0 is the expectation value at $T = 0$.

in the quantum field theory, we subtract the expectation value for $T = 0$, $\langle 0|T^{\mu\nu}|0\rangle$, to normalize it. Since the four fermion interaction is not renormalizable, we consider the spacetime dimension less than 4 to regularize the theory. Here we regard the theory for $D = 4 - 2\epsilon$ with ϵ sufficiently small positive as a regularization of the one in four dimensions. There is a correspondence between the positive regularization parameter ϵ and the cut-off scale Λ of the theory [10],

$$\frac{1}{\epsilon} - \gamma + \ln 4\pi + 1 \leftrightarrow \ln \left(\frac{\Lambda}{\mu} \right)^2, \quad (20)$$

where μ is a renormalization scale.

To obtain the expectation value of σ which corresponds to the dynamical fermion mass we evaluate the effective potential $V(\sigma)$ of the four-fermion interaction model at finite temperature. In arbitrary spacetime dimensions $2 \leq D < 4$ it is given by [15]

$$V(\sigma) = \frac{\sigma^2}{2\lambda} - \frac{1}{\beta} \frac{2tr1}{(4\pi)^{(D-1)/2} \Gamma\left(\frac{D-1}{2}\right)} \times \int dK K^{D-2} \left[\ln \cosh \left(\frac{\beta}{2} \sqrt{K^2 + \sigma^2} \right) - \ln \cosh \left(\frac{\beta}{2} K \right) \right], \quad (21)$$

at the large N limit. The expectation value of σ is determined by observing the minimum of the effective potential. In the present paper we are interested in the influence of the chiral symmetry breaking. Hence we take the coupling λ larger than the critical one and numerically calculate the behavior of σ as is shown in Fig. 1. The dynamical fermion mass is generated below a critical temperature, T_c . Details of the generated fermion mass is discussed in Ref. [15] for $2 \leq D < 4$.

Inserting the expectation value of σ into the Eqs. (18) and (19) we obtain the energy-momentum tensor of the four-fermion interaction model at finite T .

For the case of chiral symmetry restoration (i.e. $T < T_c$) the fermion is massless. We get the traceless energy-momentum tensor

$$T_{\alpha\beta} = \text{diag}(\rho, \rho/(D-1), \rho/(D-1), \rho/(D-1), \dots), \quad (22)$$

where the density ρ is introduced by

$$\rho = \frac{Ntr1}{(4\pi)^{(D-1)/2} \Gamma\left(\frac{D-1}{2}\right)} \times \int_0^\infty dK \frac{K^D}{\sqrt{K^2 + \sigma^2}} \left\{ 1 - \tanh \left(\frac{\beta}{2} \sqrt{K^2 + \sigma^2} \right) \right\}, \quad (23)$$

which is always positive by numerical calculations and formally satisfies the photon like equation of state at high temperature, $w = p/\rho = 1/(D-1)$, with $D = 4 - 2\epsilon$. The behavior for $w = p/\rho = 1/(D-1)$ is very reasonable because $\sigma = 0$ means the massless field model! So its equation of state must be similar to the photon's.

For the case of chiral symmetry breaking with σ nonzero, we have

$$T_{\alpha\beta} = \rho_0 \eta_{\alpha\beta} + \text{diag}(\rho, \rho/(D-1), \rho/(D-1), \dots) + \text{diag}(co(\sigma), 0, 0, \dots), \quad (24)$$

where the small correction

$$co(\sigma) = \frac{Ntr1}{(4\pi)^{(D-1)/2} \Gamma\left(\frac{D-1}{2}\right)} \times \int_0^\infty dK \frac{K^{D-2} \sigma^2}{\sqrt{K^2 + \sigma^2}} \times \left\{ 1 - \tanh \left(\frac{\beta}{2} \sqrt{K^2 + \sigma^2} \right) \right\}, \quad (25)$$

is also positive and the energy-momentum tensor elements also satisfy the EOS for the dark energy with the background of temperature dependent cosmological constant $\rho_0 = N[\sigma(T=0)^2 - \sigma(T)^2]/(2\lambda)$.

The behavior of the energy-momentum tensor is illustrated in Figs. 2 and 3. As is explained above, zero-point of $\langle T^{\mu\nu} \rangle$ can not be determined in the quantum field theory. We normalize it which satisfies $\langle T^{\mu\nu} \rangle = 0$ at $T = 0$. As is clearly seen, the negative pressure is realized for a low temperature region. The lines turn sharply at the critical temperature T_c .

In the following section we turn to the the classical cosmological tests of EOS for our model.

IV. DYNAMICAL EVOLUTION IN THE FRW COSMOLOGY

In addition to the Friedmann equation we have the important equation to satisfy when we solve the evolution




FIG. 2: Behavior of the energy-momentum tensor T_{00} for $\lambda = -20\sigma_0^{2-D}$ as a function of T .




FIG. 3: Behavior of the energy-momentum tensor T_{ii} for $\lambda = -20\sigma_0^{2-D}$ as a function of T .

of the universe. It is the conservation for the energy-momentum tensor of the cosmic fluid given by

$$\dot{\rho}_i + 3H(p_i + \rho_i) = 0, \quad (26)$$

where subscript i describes each species of the fluids in the considering universe. Here we mainly discuss the evolution of the universe which is dominated by the dark energy of our model.

Since the recent observational data strongly suggest that our universe is spatially flat [16, 17, 18], that is $k = 0$ in Eq.(1), we will also investigate only this case here. In this framework we mainly discuss several classical observational tests [19] for this mysterious dark energy problem.

A dimensionless quantity is more useful to discuss physical phenomena of our model. In the standard cosmology, for convenience, the density fraction of the world is expressed in terms of a critical density which is defined by the total energy density, i.e. $\rho_c = 3H_0^2/8\pi G$. When we treat the flat, homogeneous and isotropic universe, the comoving coordinates is convenient to describe the dynamical evolution of it. In this treatment the coordinates at any place do not vary with the evolution of the universe. It is described by a scale factor $a(t)$ that is a function of only cosmic time in the homogeneous and

isotropic universe. If we find the relationship between the scale factor and the cosmic time, then the evolution of the universe is clearly understood. For example, we can directly deduce the Hubble parameter and the deceleration parameters as a function of time, the dependence of the luminosity distance on redshift and so on. By investigating its dynamical evolution in the dark energy dominant era, we compare the observable quantities derived in our model with the experimental data.

We first consider a general case where equation of state is written by

$$p = w\rho, \quad (27)$$

The parameter w depend on the spacetime dimension D and this dependence is assumed to be extremely small for low temperature region. By using the continuity equation (26) and the Eq.(27), we easily obtain the evolution of the energy density ρ which obeys

$$\rho = \rho_0 a^{-n}, \quad (28)$$

where ρ_0 is the present energy density of the universe and n is given by $n = 3(1 + w)$. Substituting Eq.(28) in the Friedmann equation, we get

$$H^2 = H_0^2 [\Omega_t a^{-n} + \Omega_M a^{-D+1} + \Omega_R a^{-D}], \quad (29)$$

where H_0 is the Hubble parameter today and Ω_a is the density parameter defined by $\Omega_a = \rho_{a0}/\rho_c$. Since we consider the flat universe, the density parameters must satisfy the relation, $\sum_i \Omega_i = 1$.

It is hard to solve the Friedmann equation (29) in general situations, because we do not know the EOS for the dark energy. From Eq.(29), however, we can easily obtain the evolution of the scale factor. If we consider the dark energy dominant situation, the scale factor behaves as

$$a(t) = \left(\frac{\sqrt{n^2 H_0^2 \Omega_t}}{2} t \right)^{2/n}, \quad \text{for } n > 0$$

$$a(t) \propto \exp \left(\sqrt{H_0^2 \Omega_t} t \right). \quad \text{for } n = 0 \quad (30)$$

The latter solution shows the same behavior with the inflationally expansion when the energy scale is sufficiently high. We may use it to describe the accelerating expansion with a cosmological constant in the low energy scale, as observed today.

The dynamical evolution factor is just the function of EOS which is shown in Fig. 4. Comparing the lines in the figure, we can see that for the smaller value of n the scale factor $a(t)$ increases faster.

V. THE DECELERATION PARAMETER

Following the fact that our universe is expanding, one may ask a question that the expansion of our universe is




FIG. 4: This figure shows the scale factor of Eq.(30) as a function of cosmic time. Here we use the following values $\Omega_t = 0.75$, $H_0 = 71 \text{ km/s} \cdot \text{Mpc}$.

accelerating or decelerating now, or how fast it expands? A parameter to describe the evolution of the universe is not only the Hubble parameter but also another important parameter, so called the deceleration parameter q which is defined by

$$q = -\frac{a\ddot{a}}{\dot{a}^2}. \quad (31)$$

The parameter q is dimensionless parameter which describes the rate for slowing the expansion. By using Eq.(30) we obtain

$$q = \frac{1}{2}(n-2) = \frac{1}{2}(3w+1). \quad (32)$$

As a first consideration we assume that the state equation parameter w is constant in the dark energy dominant era.

In the four-fermion interaction model we show the evolution of the deceleration parameter in Fig. 5. As is clearly seen in Fig. 5 w is not a constant. The evolutions of w depend on both the temperature and the regularization parameter ϵ which is related to the cut-off scale of the theory. However, we think that the constant w is a good approximation when we consider the present evolution of our universe because our universe is bathed in enough low temperature now. Generally speaking, the deceleration parameter q is a function of both the present density Ω_i and w . Furthermore, it is also a function of the parameter ϵ because we consider a non-renormalizable model as a low energy effective theory. Figure 5 shows that the relationship between the parameter w and the regularization parameter ϵ . The present w (low temperature region) seems almost a monotone function of the parameter. The parameter w smoothly falls down and the curves become flat as T decreases. At the lower temperature limit w approaches the cosmological constant limit. It shows the exponential expansion of the present universe. As is shown in Eq. (22) w becomes $1/(D-1)$ at the high temperature limit. It is a property of the massless fermion. Therefore the exponential expansion only appears at the low temperature region in our model without any fine tunings. It is a very important property




FIG. 5: The evolution of the state equation parameter w with respect to T/σ_0 . We set $\lambda = -20\sigma_0^{2-D}$.

to a dark energy candidate. Slopes of each lines in Fig.5 are depend on the coupling λ . For larger coupling λ the slope becomes steeper. However, the behaviors of w does not change qualitatively as λ varies.

From the recent SN Ia and Cosmic Microwave Background observations [5] a cosmological constant-like dark energy may exist today which accelerate the expansion of the universe. It suggests that the decelerating parameter lies in a range, $-1 < q < 0$. This observational constraint is naturally satisfied in the four-fermion interaction model. Recently there are many dark energy models of simple scalar fields like the quintessence [6], however, they all need fine tuning of some parameters to certain extents. In the present model, on the other hand, fine tuning is not necessary. In other words the model has some properties which are realized in the quintessence scenario without any cosmological constant-like terms. Certainly we need more SN Ia observational data to confirm the important constraint, which is just the main mission of the near future SNAP project.

VI. CONCLUDING REMARKS

We investigated the cosmological effects of the vacuum expectation value of energy momentum tensor in a four fermion interaction model. According to the imaginary time formalism we derive the explicit form of the energy momentum tensor. The energy-momentum tensor depends on the temperature and the dynamically generated fermion mass. The fermion field is massless and has the similar property as the radiation field above the critical temperature. We consider that the temperature of the present universe is much less than the critical one and the chiral symmetry is broken down dynamically. Thus the composite operator of the fermion and the anti-fermion develops a non-vanishing vacuum expectation value in the present stage of the universe. The vacuum expectation value plays a role of the cosmological constant-like.

Applying it to models of the universe without an explicit cosmological constant term, we discussed the de-

pendence of the scale factor $a(t)$ on the state equation parameter w and evaluate the deceleration parameter q . We found that the behavior of the EOS is dramatically changed at low temperature. A dark energy property was naturally obtained in the four-fermion interaction model. In the present analysis the fermion mass scale can be at the scale of neutrino, QCD, electroweak symmetry breaking, extra-dimension [20] and so on. We have had a loosely constrained parameter range for the cosmological constant [21]. For more concrete analysis we must apply the results to some cosmological tests with comparisons to observational data, which are under our investigations.

Our model is only a prototype one, but we believe that the results are worth discussing further about the real world. By observational data analysis we know the dark energy is almost smoothly distributed [22], which implies a kind of very weak interaction that we are not very clear about yet. Here we just employ the simple four-fermion interaction model to give the global energy momentum tensor which depends on temperature and some model parameters. It might be too simple, however we found interesting results. As a next step, it is very important to apply our analysis to other well-defined and more realistic models. One of the interesting models is SUSY NJL model [23, 24]. For the SUSY NJL model the effective action is written as

$$\Gamma[\sigma] = \int d^4x [\text{Tr} \ln(i \not{\partial} - \sigma) - 2\text{Tr} \ln(\partial^2 - \sigma^2) + N\sigma^2/(2\lambda)]. \quad (33)$$

According to the supersymmetry, contributions from the fermion and boson loops cancel out each other exactly at $T = 0$. Since the thermal effect restore the chiral

symmetry, dynamical symmetry breaking does not take place even at finite temperature. If we introduce a soft SUSY breaking term such as $\Delta\phi^*\phi$ and/or $\Delta\phi^{c*}\phi^c$ in the action [25], then the contribution from the boson loop is modified. The composite operator $\bar{\psi}\psi$ can develop a non-vanishing vacuum expectation value which satisfies the dark energy property to some range. The scalar mass depends on the coupling parameter. When the coupling parameter is large enough, $\Delta > 2\sigma$, the scalar mass $m = \sqrt{(\sigma^2 - \Delta)}$ becomes heavier than the dynamical fermion mass. We can choose the coupling parameter Δ to realize the appropriate scale for the cosmological constant and the negative pressure of the present universe accordingly.

So far the origin of the dark energy in the current cosmological models is still under discussions with many theoretical speculations. It is accommodated in the standard Friedmann cosmology based on the relativistic gravitational theory with a cosmological constant-like term. It can also be interpreted in extended gravity scenarios and corresponding cosmological models, like the brane cosmologies. That attracts us to further research works continuously.

Acknowledgments

The paper is started in X. H. Meng's academic visit at Hiroshima University supported from JSPS INVITATION Fellowship. The authors have also benefited a lot from discussions with many people, especially, D. Kimura, G. Senjanovich, X. P. Wu, K. Yamamoto and Z. H. Zhu. This work is also supported partly by grant of No.NK-BRSF G19990753 from Ministry of Science and Technology of China.

-
- [1] See for example, S. Perlmutter, M. S. Turner and M. White, Phys. Rev. Lett. **83**, 670 (1999).
 - [2] P. Bernadis, *et al*, Nature **404**, 955 (2000); S. Hanany, *et al*, Astrophys. J. Lett. **545**, L5 (2000); A. Balbi, *et al*, Astrophys. J. Lett. **545**, L1 (2000); M. Collers, astro-ph/0305051; D. Spergel, *et al*, astro-ph/0302209; C. Bennet, *et al*, astro-ph/0302207 for example.
 - [3] S. Perlmutter, *et al*, Nature **391**, 51 (1998); J. Tonry, *et al*, astro-ph/0305008.
 - [4] For example see, M. R. Nolte, *et al*, astro-ph/0305097 and references therein
 - [5] S. Perlmutter, *et al*, Nature **404**, 955 (2000); Astrophys. J. **517**, 565 (1999); A. Riess, *et al*, Astrophys. J. **116**, 1009 (1998); *ibid*, **560**, 49 (2001); P. Peebles, astro-ph/0208037; astro-ph/0207347 and references therein.
 - [6] For example see, P. Peebles and B. Ratra, Astrophys. J. Lett. **325**, L17 (1998); C. Wetterich, Nucl. Phys. B **302**, 668 (1998); L. A. Boyle, *et al*, astro-ph/0105318; C. A. Picon, *et al*, astro-ph/0004134.
 - [7] J. I. Kapusta, *Finite-Temperature Field Theory* (Cambridge University Press, 1989); L. Bellac, *Thermal Field Theory* (Cambridge University Press, 1996); A. Das, *Finite Temperature Field Theory* (World Scientific, 1997).
 - [8] T. Inagaki, T. Muta and S. D. Odintsov, Mod. Phys. Lett. A **8**, 2117 (1993); T. Inagaki, S. Mukaigawa and T. Muta, Phys. Rev. **D52**, 4267 (1995); K. Ishikawa, T. Inagaki and T. Muta, Mod. Phys. Lett. A **11**, 939 (1996); T. Inagaki, Int. J. Mod. Phys. A **11**, 4561 (1996); T. Inagaki and K. Ishikawa, Phys. Rev. **D56**, 5097 (1997).
 - [9] N. Rius and V. Sanz, Phys. Rev. **D64**, 075006 (2001).
 - [10] T. Inagaki, T. Muta and S. D. Odintsov, Prog. Theor. Phys. Suppl. **127**, 93 (1997).
 - [11] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
 - [12] D. J. Gross and A. Neveu, Phys. Rev. **D10**, 3235 (1974).

- [13] N. D. Birrell and P. C. W. Davies, *Quantum Field Theory in Curved Space* (Cambridge University Press, 1982); S. A. Fulling, *Aspects of Quantum Field Theory in Curved Space-Time* (Cambridge University Press, 1989); R. M. Wald, *Quantum Field Theory in Curved Space-time and Black-Hole Thermodynamics* (The University of Chicago Press, 1994).
- [14] L. Dolan and R. Jackiw Phys. Rev. **D9**, 3320 (1974), S. Weinberg, Phys. Rev. **D9**, 3357 (1974).
- [15] T. Inagaki, T. Kouno and T. Muta, Int. J. Mod. Phys. A **10**, 2241 (1995).
- [16] C. Pryke, astro-ph/0104490.
- [17] C. B. Netterfield, *et al*, astro-ph/0104460.
- [18] R. Stompor, astro-ph/0105062.
- [19] X. Meng, *et al*, Int. J. Mod. Phys. D **3**, 369(2003); hep-ph/0210096.
- [20] For example see, H. Abe, K. Fukazawa and T. Inagaki, Prog. Theor. Phys. **107**, 1047 (2002); H. Abe and T. Inagaki, Phys. Rev. **D 66**, 085001 (2002).
- [21] S. Weinberg, Phys. Rev. Lett. **59**, 2697 (1987); Rev. Mod. Phys. **61**, 1 (1989).
- [22] P. P. Avelino, J. P. M. de Carvalho and C. J. A. P. Martins, Phys. Rev. **D64**, 063505 (2001).
- [23] W. Buchmuller and S. T. Love, Nucl. Phys. B **204**, 213 (1982).
- [24] I. L. Buchbinder, T. Inagaki and S. D. Odintsov, Mod. Phys. Lett. A **12**, 2271 (1997); T. Inagaki, S. D. Odintsov and Y. I. Shil'nov, Int. J. Mod. Phys. A **14**, 481 (1999).
- [25] J. Hashida, T. Muta and K. Ohkura, Phys. Lett. B **451**, 343 (1999); J. Hashida, T. Muta and K. Ohkura, Mod. Phys. Lett. A **13**, 1235 (1998).